## 2023 Recommended Immunizations for Children from Birth Through 6 Years Olds



## **FOOTNOTES**





Administering a third dose at age 6 months depends on the brand of Hib or rotavirus vaccine used for previous dose. COVID-19\*\* Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

Flut Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.

HepA<sup>‡</sup> n

Two doses of Hep A vaccine are needed for lasting protection.
The 2 doses should be given

between age 12 and 23 months. Both doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.

## ADDITIONAL INFORMATION

1. If your child misses a shot recommended for their age, talk to your child's doctor as soon as possible to see when the missed shot can be given.

2. If your child has any medical conditions that put them at risk for infection (e.g., sickle cell, HIV infection, cochlear implants) or is traveling outside the United States, talk to your child's doctor about additional vaccines that they may need.

Talk with your child's doctor if you have questions about any shot recommended for your child.

U.S. Department of Health and Human Services

Centers for Disease Control and Prevention

FOR MORE INFORMATION

Call toll-free: 1-800-CDC-INFO (1-800-232-4636)

Or visit: cdc.gov/vaccines/parents

This content was adapted by Pfizer from the CDC's 2023 childhood immunization schedule.

§This is a resource intended for parents or caregivers; there is a more detailed healthcare professional version of the schedule available on the CDC website.

PP-PRV-USA-0299 © 2023 Pfizer Inc. All rights reserved. February 2023

